

## TUBERCULOSIS INFECTION ON DYSFUNCTIONAL FARMS

1Masharipova Sh. S.

2 Masharipov S. M.

1Urgench Branch of Tashkent Medical Academy, Urgench, Uzbekistan

2Alfraganus University, Tashkent, Uzbekistan

### Abstract:

In 2022, tuberculosis claimed the lives of a total of 1.3 million people (including 167,000 people infected with HIV). Worldwide, tuberculosis is the second leading cause of death from infectious diseases after COVID-19 (mortality rates are higher than from HIV infection and AIDS). The urgency of the problem of tuberculosis incidence is associated with the spread of drug-resistant strains of the causative agent of tuberculosis, high susceptibility to the disease of unvaccinated persons, possible chronization of the process with late medical treatment and refusal of etiotropic treatment. It is known that sick animals and small cattle cause human tuberculosis infection. The symptoms of tuberculosis in livestock have not been sufficiently studied in the literature, which significantly complicates the clinical recognition of tuberculosis in animals. There are no effective methods of diagnosis and specific prevention of the disease.

**Keywords:** Mantoux test, BCG vaccination, epidemic, focus of infection, epizootic, temperature, causative agent of tuberculosis, population, pets.

### Introduction

Tuberculosis is a socially significant and especially dangerous infectious disease. The main source of infection is a person with tuberculosis, less often cattle, camels, pigs, birds, and other animals. According to WHO, one third of the world's population is infected with tuberculosis. 8 million new cases of tuberculosis and 3 million deaths from it are registered annually in the world, including 884,000 children under the age of 15. The total number of tuberculosis patients in the world today reaches 60 million, and about a third of the world's population is infected. According to WHO forecasts, 90 million new cases of the disease are expected in the current millennium. An analysis of the epidemiological situation of tuberculosis in the world over the past decade has shown that forecasts of the elimination of tuberculosis as a common disease have not been realized. In 1993, WHO declared tuberculosis a global scourge for humanity. Every year, the number of tuberculosis patients increases by 8-10 million and of them 3-4 million die from tuberculosis, women make up about 1 million, and children – more than three hundred thousand.

Tuberculosis is a widespread infectious disease of humans and animals in the world caused by various types of Mycobacterium tuberculosis complex Mycobacterium, or otherwise Koch's sticks, capable of living inside cells. Tuberculosis usually affects the lungs, less often affecting



other organs and systems. Mycobacterium tuberculosis is transmitted by airborne droplets during conversation, coughing and sneezing of the patient. Most often, after infection with mycobacteria, the disease proceeds in an asymptomatic, latent form (tubification), but about one in ten cases of latent infection eventually turns into an active form. For a person, the disease is socially dependent.

### **The Purpose of the Study**

He will study tuberculosis infection on disadvantaged farms from animals.

### **Materials and Methods of Research**

In accordance with the "Law on Veterinary Medicine" of the Republic of Uzbekistan, it is necessary to ensure the protection of the population from infectious diseases common to humans and animals. Given that an animal is the cause of human infection, great attention should be paid to epizootic monitoring, including those designed to track the epizootic situation in order to optimize prognostic conclusions and management decisions to reduce risks to the population. But, despite the success in developing a practical and theoretical basis for monitoring the epizootic situation of infectious diseases, it is quite obvious that it needs to be improved in case of significant diseases for specific territories.

Tuberculosis poses a serious danger, despite the progress in public health: a third of the world's population is infected. And only together can this disease be overcome.

### **Research Results**

Tuberculosis is a chronic infectious disease characterized by the formation of tuberculosis in parenchymal organs, intestines and other tissues. The disease is caused by the causative agent of tuberculosis Mycobacterium tuberculosis. There are several main types of pathogen pathogenic to humans and animals. Modern scientific data have established that the causative agent of tuberculosis can affect animals not only of the species to which it has adapted, but also many other species.

In animals, 5 pathogenic mycobacterium species have epizootic significance: 1) M.bovis is the causative agent of tuberculosis in cattle, pathogenic for domestic and wild ruminants, humans; 2) M.tuberculosis is the causative agent of human tuberculosis, pathogenic for humans, primates, dogs, parrots; 3) M.avium is the causative agent of avian tuberculosis, pathogenic for birds, pigs, cats, dogs; 4) M.microti is the causative agent of tuberculosis in mice, mouse-like rodents; 5) M.piscium is the causative agent of tuberculosis in cold-blooded.

Thus, a human-type pathogen, except for humans, can infect dogs, cats, pigs, monkeys, parrots, and guinea pigs from laboratory animals. The causative agent of bovine tuberculosis, in turn, in addition to cattle, can also infect humans, cats, pigs, goats, sheep, dogs and hares. The causative agent of avian tuberculosis affects pigs, horses, cats and, as has been proven recently, quite often humans. It is not possible to determine the exact number of unproductive pets, but if we assume that every third Uzbek family contains a cat, and every fifth family contains a dog, then, according to independent experts, there may be at least 15.9 million dogs and cats in



Uzbekistan. At the same time, the proportion of purebred dogs and cats does not exceed 2%. Based on the data obtained, it is assumed that Central Asia occupies the fifth place in the world and the leading position in Europe in terms of the number of cats and dogs. Thus, small pets are the most numerous and common. The peculiarity of the habitat of unproductive animals of cats and dogs is that their habitat is populated areas and directly human habitation. This factor can significantly negatively affect the hygiene of the environment: the lack of special places for walking dogs and cats, uncontrolled charity and the availability of food waste contribute to the growth of the number of stray animals in cities and towns. In addition, this situation attracts wild and synanthropic animals to human habitation, which can, in turn, lead to an unfavorable epizootic situation.

Also, the habitation of animals (cats and dogs) among humans contributes to the spread of infectious and invasive diseases of humans and other domestic animals. So, for example, a dog is a source of many infectious and parasitic diseases transmitted to humans. Unfortunately, in our country there is no statistical database on the number of small domestic (unproductive) animals, so it is not possible to track the epizootological geography of infectious diseases of dogs and cats. Studies of the features of epizootic processes in various regions are of extreme importance in the development of a set of antiepidemic measures. Despite the achievements of veterinary science and practice in the prevention and treatment of infectious animal diseases, these diseases continue to damage the economies of various countries. Due to the development of globalization processes and the deterioration of the environmental situation, close contacts between humans and various animal species occur much more often than before.

### **Discussion**

Therefore, there is evidence that new diseases may appear in the near future that can cause even more damage. In particular, this is due to the fact that some animals that did not previously live in close proximity to humans have been domesticated, and the modern food production system is imperfect and can lead to animal epidemics that can threaten human health and life. Climate change has led to changes in the habitats of migratory insects and birds, which are carriers of pathogens of various diseases. In addition, changes in the habitat of humans and animals also lead to changes in the resistance and pathogenesis of pathogens of infectious diseases.

According to the data provided by the World Health Organization, there are approximately nosological forms of diseases in the world that are common to animals and humans. These diseases include anthrax, rabies, tuberculosis, brucellosis, glanders, tick-borne encephalitis, leptospirosis, foot-and-mouth disease, actinomycosis, Cu fever, echinococcosis, diphyllbothriosis, salmonellosis, etc. Pets are ubiquitous. It is worth considering that the number of these animals directly depends on the degree of development of urbanized territories. The population density of dogs and cats can reach huge numbers. Thus, according to WHO, the global dog population was estimated at more than 0.5 billion individuals. And the number of cats in the world does not exclude 1 billion animals. All sick animals (regardless of the type of pathogen) pose a danger to human health. In turn, a person with tuberculosis is a source of infectious agent for animals, from which other healthy people can become infected. Many cases



of the incidence of human-type tuberculosis in monkeys of one of the menageries have been described. All of the above indicates the need for a comprehensive fight against tuberculosis in both humans and all tuberculosis-susceptible animal species. In recent years, simultaneously with the growth of the well-being of the urban population, there has been a tendency to increase the number of small pets up to critical indicators. At the same time, there are statistics that the number of dogs and cats has increased by 1.4 times in recent decades, and the population by 1.2 times.

Most multiply resistant bacteria are characterized by reduced virulence – the ability to spread and cause disease in the host organism. New strains of tuberculosis combined with climate-related crop failures will lead to mass human migration and infection of small animals. For thousands of years, tuberculosis has continued to be one of the most difficult problems of human and animal infectious pathology, causing significant economic damage to livestock and posing a real danger of human infection.

Animals transported across the border of the Republic of Uzbekistan must be kept in quarantine for at least 21 days, during which thermometry and diagnostic tests of animals for brucellosis and tuberculosis are carried out.

Currently, in many countries there is an increase in the incidence of tuberculosis in people, especially children. Tuberculosis occupies a special place among infectious animal diseases. It is peculiar in that for many years it can occur in a latent form, without showing clinical signs of the disease, without affecting the productivity and vital activity of animals. Dogs and cats are an integral part of megacities, provincial towns, villages and villages, where they are in close contact with humans and other animal species.

### **Conclusion**

Thus, studying the role and place of infectious diseases in the general structure of pathology of domestic animals, clarifying the features of their epizootic manifestation in the urban ecosystem, as well as improving the epizootic surveillance system for infectious diseases of domestic carnivores is a very urgent task, the solution of which requires urgent solution. The emergence of antibiotic-resistant tuberculosis has become a special danger, unlike the threat from the so-called superbugs and much more serious.

### **References**

1. Юсупов Ш.Р., Аскарова Р.И., Машарипова Ш.С., Якубова У.Б. АНАЛИЗ ФАКТОРОВ РИСКА, ВЛИЯЮЩИХ НА РАЗВИТИЕ ТУБЕРКУЛЕЗА У ДЕТЕЙ В ХОРЕЗМСКОЙ ОБЛАСТИ / Наука, техника и образование 2019. № 8 (61) стр. 66-73.
2. Юсупов Ш.Р. Туберкулез в Хорезмской области/International Scientific Review of the Problems of Natural Sciences and Medicine / Boston. USA. 2020 г. апрель – стр. 16-23.
3. Машарипова Ш.С., Матякубова О.У ТЕЧЕНИЕ ВИЧ/СПИД ИНФЕКЦИИ У БОЛЬНЫХ ТУБЕРКУЛЕЗОМ ЛЕГКИХ / European science. 2020 г. № 3 (52) – стр. 110-113.



4. Аскарлова Р.И., Юсупов Ш.Р., Машарипова Ш.С., Машарипова Х.К. Эпидемиология легочного туберкулеза / EUROPEAN RESEARCH: INNOVATION IN SCIENCE, EDUCATION AND TECHNOLOGY LVII International correspondence scientific and practical conference. 2019 г. – стр. 96-100.
5. Машарипов С.М., Юсупов Ш.Р., Машарипова Ш.С., Матякубова О.У. КЛИНИЧЕСКОЕ ТЕЧЕНИЕ ТУБЕРКУЛЕЗА У БОЛЬНЫХ ГЕПАТИТОМ В / Вестник ТМА.uz. 2023 г. № 3 (2) - стр. 155-157.
6. Masharipov S. M. et al. CHICKENPOX DISEASE AND ITS SPECIFIC PREVENTION //Лучшие интеллектуальные исследования. – 2024. – Т. 20. – №. 3. – С. 131-138.
7. Sobir M. TUBERCULOSIS AND ITS COURSE IN PATIENTS WITH HEPATITIS B. – 2023.
8. Sobir M. S. S. M. IMMUNE STATUS OF ADULTS AND CHILDREN WITH AN ALLERGIC BACKGROUND DIAGNOSED WITH ENTEROBIOSIS. – 2023.
9. Masharipov S. M. et al. THE SPREAD OF ENTEROBIOSIS IN THE CITY OF URGENCH. – Сборник тезисов международной научно-практической конференции «Современные тенденции развития инфектологии, медицинской паразитологии, эпидемиологии и микробиологии», 2023.
10. Машарипова Ш. Specific pathomorphological changes in the course of lymphocytic leukosis in children //Современник аспекты паразитологии и актуальные проблемы кишечных инфекций. – 2024. – Т. 1. – №. 1. – С. 25-25.
11. Машарипова Ш. Frequently ill children in the khorezm region //Современник аспекты паразитологии и актуальные проблемы кишечных инфекций. – 2024. – Т. 1. – №. 1. – С. 26-26.
12. Машарипова Ш. С., Ёкубова У. Б. УХОД ЗА ИНФЕКЦИОННЫМ БОЛЬНЫМ И ЕГО ПИТАНИЕ //Современная наука: актуальные проблемы и пути их решения. – 2017. – №. 2. – С. 81-82.
13. Хударгенова Д. Р. и др. ЦЕЛЕСООБРАЗНОСТЬ ОТБОРА ЖИВОТНЫХ-ПРОДУЦЕНТОВ ПРИ ПОЛУЧЕНИИ ГИПЕРИММУННЫХ СЫВОРОТОК, ПРИМЕНЯЕМЫХ В СУДЕБНО-МЕДИЦИНСКИХ ЛАБОРАТОРИЯХ. – 2023.
14. Ibadullayeva S. S. et al. FEATURES OF INFECTIOUS MONONUCLEOSIS IN CHILDREN //Scientific Impulse. – 2023. – Т. 2. – №. 16. – С. 1165-1171.
15. Машарипова Ш. С. О ‘РКА ARTERIYALARINING QANDLI DIABET TA’SIRIDA MORFOLOGIK TUZILISHI //Журнал кардиореспираторных исследований. – 2022. – Т. 3. – №. 1.
16. Sadullaev, S. E., et al. "PREVALENCE OF DIARRHEAL DISEASES IN THE REPUBLIC OF UZBEKISTAN." International Journal of Education, Social Science & Humanities 12.3 (2024): 356-363.
17. Sadullaev S. E., Sh M. D. THE COURSE OF CORONAVIRUS AGAINST THE BACKGROUND OF CHRONIC HEPATITIS Masharipova Sh. S Masharipov S //Научный импульс. – 2023. – С. 78.



---

18.Masharipova Sh.S, & Masharipov S.M. (2024). CHARACTERISTIC CHANGES IN ARTERIAL VESSELS IN TYPE 2 DIABETES MELLITUS. *Scientific Impulse*, 2(20), 374–378.

19.Masharipova, S., Mirsharapov, U., Masharipov, S., & Mahsripov, A. (2021). Living Vessel Morphological Structure.

